

Homology operations on homology of quandles

M. Niebrzydowski

University of Louisiana at Lafayette, Department of Mathematics, 1403 Johnston St., Lafayette, LA 70504-1010

J.H. Przytycki^{*}

The George Washington University, Department of Mathematics, Monroe Hall, Room 240, 2115 G St. NW, Washington, D.C. 20052

Abstract

We introduce the concept of the quandle partial derivatives, and use them to define extreme chains that yield homological operations. We apply this to a large class of finite and infinite quandles to show, in particular, that they have nontrivial elements in the third and fourth quandle homology.

Key words: Dihedral quandle, Alexander quandle, Rack, Quandle homology, Homological operation, Quandle partial differential equation
1991 MSC: 55N35, 18G60, 57M25

Quandles are algebraic structures introduced by David Joyce in his 1979 Ph.D. thesis [14] as a powerful tool for classifying knots (compare [15] and S. Matveev [18]). Rack homology and homotopy theory were first defined and studied in [11], and a modification to quandle homology theory was given in [3] to define knot invariants in a state-sum form (so-called cocycle knot invariants). In this paper, we consider various homological operations on homology of quandles. We introduce the notion of quandle partial derivatives, and extreme chains on which appropriate partial derivatives vanish. We also consider the degree one homology operations created using elements of the quandle satisfying the so-called k -condition.

^{*} Corresponding author.

Email addresses: mniebrz@gmail.com (M. Niebrzydowski), przytyck@gwu.edu (J.H. Przytycki).

1 Definitions and preliminary facts

Definition 1 A *quandle*, X , is a set with a binary operation $(a, b) \mapsto a * b$ such that

- (1) For any $a \in X$, $a * a = a$.
- (2) For any $a, b \in X$, there is a unique $c \in X$ such that $a = c * b$.
- (3) For any $a, b, c \in X$, $(a * b) * c = (a * c) * (b * c)$ (right distributivity).

Note that the second condition can be replaced with the following requirement: the operation $*_b: Q \rightarrow Q$, defined by $*_b(x) = x * b$, is a bijection.

Definition 2 A *rack* is a set with a binary operation that satisfies (2) and (3).

We use a standard convention for products in non-associative algebras, called the left normed convention, that is, whenever parentheses are omitted in a product of elements $a_1, a_2, \dots, a_n$ of Q then

$$a_1 * a_2 * \dots * a_n = ((\dots((a_1 * a_2) * a_3) * \dots) * a_{n-1}) * a_n$$

(left association). For example, $a * b * c = (a * b) * c$.

According to [10], the earliest discussion on racks is in the correspondence between J. H. Conway and G. Wraith, who studied racks in the context of the conjugacy operation in a group. They regarded a rack as the wreckage of a group left behind after the group operation is discarded and only the notion of conjugacy remains.

There are several basic constructions of quandles from groups and $\mathbb{Z}[t^\pm]$ -modules. We list them below stressing the functorial character of their definition. In particular, the category of quandles has quandles as objects and quandle homomorphisms as morphisms.

Definition 3

- (i) There is a functor T from the category of abelian groups to the category of quandles, such that for a group G , $T(G)$ is a quandle with the same underlying set as G and with quandle operation $*$ given by $a * b = 2b - a$. For a group homomorphism $f : G \rightarrow H$, we define $T(f) = f$ as a function on the set. We check that $T(f)$ is a quandle homomorphism: $T(f)(a * b) = f(a * b) = f(2b - a) = 2f(b) - f(a) = f(a) * f(b) = T(f)(a) * T(f)(b)$. This construction was first considered by M. Takasaki [24], so we call this functor the *Takasaki functor*.

- (ii) There is a functor from the category of groups into the category of quandles in which $a * b = b^{-1}ab$. This functor is called the *conjugacy functor*.
 - (iii) There is a functor from the category of groups into the category of quandles, in which $a * b = ba^{-1}b$. This functor, generalizing Takasaki functor, is called the *core functor*.
 - (iv) There is a functor from the category of $\mathbb{Z}[t^{\pm 1}]$ -modules into the category of quandles in which $a * b = (1 - t)b + ta$. This functor, also generalizing Takasaki functor, is called the *Alexander functor*.
- In particular, if $f: A_1 \rightarrow A_2$ is a $\mathbb{Z}[t^{\pm 1}]$ -homomorphism, then it is also an Alexander quandle homomorphism.

We recall below the notion of rack and quandle homology. It is useful, following Kamada [16], to place it in a slightly more general setting, in which we deal with a rack and a rack-set on which the rack acts:

Definition 4

- (1) For a rack (or a quandle) X , the set Y is a *rack-set* (or *X-set*) if there is a map $*$: $Y \times X \rightarrow Y$, such that
 - (i) the map $*_x : Y \rightarrow Y$, given by $*_x(y) = y * x$, is a bijection, and
 - (ii) $(y * a) * b = (y * b) * (a * b)$.
- (2) For a given rack X and a rack-set Y , let $C_n^R(X, Y)$ be the free abelian group generated by n -tuples $(y, x_2, \dots, x_n)$, with $y \in Y$ and $x_i \in X$, $i = 2, \dots, n$; in other words, $C_n^R(X, Y) = \mathbb{Z}(Y \times X^{n-1}) = \mathbb{Z}Y \otimes \mathbb{Z}X^{\otimes n-1}$. Define a boundary homomorphism $\partial : C_n^R(X, Y) \rightarrow C_{n-1}^R(X, Y)$ by:

$$\begin{aligned} \partial(y, x_2, \dots, x_n) = & \sum_{i=2}^n (-1)^i ((y, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \\ & - (y * x_i, x_2 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n)). \end{aligned}$$

$(C_*^R(X, Y), \partial)$ is called the *rack chain complex of the pair* (X, Y) .

The homology of this chain complex is called the *homology of the pair* (X, Y) .

- (3) Assume that X is a quandle. Then we have a subchain complex of late degenerate elements, $C_n^{DD}(X, Y) \subset C_n^R(X, Y)$, generated by n -tuples $(y, x_2, \dots, x_n)$ with $x_{i+1} = x_i$ for some i . The subchain complex $(C_n^{DD}(X, Y), \partial)$ is called the *late degenerated chain complex of the pair* (X, Y) . The homology of this chain complex is called the *late degenerated homology of* (X, Y) , and the homology of the quotient chain complex $C_n^{LQ}(X, Y) = C_n^R(X, Y)/C_n^{DD}(X, Y)$ is called the *late quandle homology of the pair* (X, Y) .
- (4) If X is a quandle and Y is an X -invariant subquandle of X (that is, $y * x \in Y$ for any $y \in Y$ and $x \in X$), then we have a subchain complex $C_n^D(X, Y) \subset C_n^R(X, Y)$, generated by n -tuples $(x_1, \dots, x_n)$ with $x_1 \in Y$ and $x_{i+1} = x_i$ for some $i \in \{1, \dots, n-1\}$. The subchain complex

$(C_n^D(X, Y), \partial)$ is called the *degenerated chain complex of a quandle pair* (X, Y) .

The quotient chain complex $C_n^Q(X, Y) = C_n^R(X, Y)/C_n^D(X, Y)$ is called the *quandle chain complex of a pair* (X, Y) .

If $X = Y$ then we deal with classical quandle homology theory.

Free part of homology of finite racks or quandles ($free(H_*(X))$) was computed in [17,8] (lower bounds for Betti numbers were given in [4]).

Theorem 5 [4,17,8] *Let $\mathcal{O}$ denote the set of orbits of a rack X with respect to the action of X on itself by the right multiplication (we say that a rack is connected if it has one orbit).*

Then:

- (i) $rank H_n^R(X) = |\mathcal{O}|^n$ for a finite rack X ;
- (ii) $rank H_n^Q(X) = |\mathcal{O}|(|\mathcal{O}| - 1)^{n-1}$ for a finite quandle X ;
- (iii) $rank H_n^D(X) = |\mathcal{O}|^n - |\mathcal{O}|(|\mathcal{O}| - 1)^{n-1}$ for a finite quandle X ;
- (iv) $rank H_n^{DD}(X) = |\mathcal{O}|^n - |\mathcal{O}|^2(|\mathcal{O}| - 1)^{n-2}$ for a finite quandle X , where $H_n^{DD}(X)$ denotes the delayed degenerated homology.

If we assume that X satisfies certain mild conditions (as in Theorem 17), then Theorem 5 follows from Theorem 17 inductively.

We should stress here, that the assumption of X being finite is essential, as M. Eisermann demonstrated that fundamental quandles of nontrivial knots (so, infinite and connected quandles) have H_2^Q equal to $\mathbb{Z}$ [9].

For the dihedral quandle $R_k = T(\mathbb{Z}_n)$ (that is, the set $\{0, 1, \dots, n-1\}$ with $a * b = 2b - a$ modulo n), we have:

$$\textbf{Corollary 6} \quad free(H_n^Q(R_k)) = \begin{cases} \mathbb{Z} & \text{for } n = 1, k \text{ odd} \\ 0 & \text{for } n > 1, k \text{ odd } (R_k \text{ is connected}) \\ \mathbb{Z}^2 & \text{for } k \text{ even } (R_k \text{ has two orbits}) \end{cases}$$

Useful information concerning torsion of homology of racks and quandles was obtained in [17,19,21]. In particular, it was shown that:

Theorem 7

- (i) [17]

$$H_2^R(X) \cong H_2^Q(X) \oplus \mathbb{Z}\mathcal{O}_X,$$

$$H_3^R(X) \cong H_3^Q(X) \oplus H_2^Q(X) \oplus \mathbb{Z}\mathcal{O}_X^2.$$

In particular, $H_3^R(R_k) \cong H_3^Q(R_k) \oplus \mathbb{Z}$ for k odd.

(ii) [21] $H_3^Q(R_p) = \mathbb{Z}_p$, and $\mathbb{Z}_p \subset H_4^Q(R_p)$ for p an odd prime.

Quandle homology was generalized to twisted quandle homology in [2], and further in [1,23] using the notion of quandle algebra. Some of our results apply to these generalizations but we will not address it in this paper.

2 Homological operations obtained from k -elements

Generalizing [21], we consider the degree one homological operation related to the group homomorphism $h_a: C_n^R(X) \rightarrow C_{n+1}^R(X)$, given by $h_a(w) = (w, a)$, for any $a \in X$, and $w \in X^n$. In general, the map h_a is not a chain map, so we need to symmetrize it with respect to another map $*_a: C_n^R(X) \rightarrow C_n^R(X)$ given by $*_a(w) = w * a$, for any $w \in X^n$, or more precisely, $*_a(x_1, \dots, x_n) = (x_1, \dots, x_n) * a = (x_1 * a, \dots, x_n * a)$. For a symmetrization we need an element $a \in X$ which satisfies a k -condition, that is, for any $x \in X$ we have $x * a^k = x$ (a quandle in which every element satisfies the k -condition is called a k -quandle; in particular a 2-quandle is an involutive quandle or a kei).

For a k -element a , we consider a function

$$h'_a = h_a + *_a h_a + *_a *_a h_a + \dots + (*_a)^{k-1} h_a.$$

The basic properties of the maps $*_a$, h_a and h'_a are described in the following proposition:

Proposition 8

- (i) *For any rack X , and $a \in X$, the map $*_a: C_n^R(X) \rightarrow C_n^R(X)$ is a chain map chain homotopic to the identity.*
- (ii) *If a satisfies the k -condition, then $h'_a: C_n^R(X) \rightarrow C_{n+1}^R(X)$ is a chain map. Notice that if X is finite, then any $a \in X$ satisfies a k -condition for some k .*
- (iii) *If X is a quandle, then $h_a(C_n^D(X)) \subset C_{n+1}^D(X)$, $*_a(C_n^D(X)) \subset C_n^D(X)$. Therefore, the maps $h_a, h'_a: C_n^Q(X) \rightarrow C_{n+1}^Q(X)$, and $*_a: C_n^Q(X) \rightarrow C_n^Q(X)$ are well defined. Furthermore, $*_a$ and h'_a are chain maps, and $*_a$ is chain homotopic to identity.*
- (iv) *If a and b are in the same orbit of X , then h'_a and h'_b induce the same map on homology, that is, $(h'_a)_* = (h'_b)_*: H_n^W(X) \rightarrow H_{n+1}^W(X)$, where $W = R, Q$ or D .*

PROOF. (i) $*_a$ is a chain map, because

$$\begin{aligned}
d(*_a(x_1, \dots, x_n)) &= d(x_1 * a, \dots, x_n * a) = \\
&\sum_{i=2}^k (-1)^i ((x_1 * a, \dots, x_{i-1} * a, x_{i+1} * a, \dots, x_n * a) \\
&- (x_1 * a * (x_i * a), \dots, x_{i-1} * a * (x_i * a), x_{i+1} * a, \dots, x_n * a)) = \\
&\sum_{i=2}^k (-1)^i ((x_1 * a, \dots, x_{i-1} * a, x_{i+1} * a, \dots, x_n * a) \\
&- (x_1 * x_i * a, \dots, x_{i-1} * x_i * a, x_{i+1} * a, \dots, x_n * a)) = \\
&*_a \left(\sum_{i=2}^k (-1)^i ((x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \right. \\
&\left. - (x_1 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n)) \right) = *_a(d(x_1, \dots, x_n)).
\end{aligned}$$

The homomorphism $(-1)^{n+1}h_a: C_n^R(X) \rightarrow C_{n+1}^R(X)$ is a chain homotopy between Id and $*_a$ chain maps. Namely:

$$\begin{aligned}
d((-1)^{n+1}h_a(x_1, \dots, x_n)) &= d((-1)^{n+1}(x_1, \dots, x_n, a)) = \\
(-1)^{n+1}((d(x_1, \dots, x_n), a) &+ (-1)^{n+1}((x_1, \dots, x_n) - (x_1 * a, \dots, x_n * a))) = \\
-(-1)^n h_a(d(x_1, \dots, x_n)) &+ (Id - *_a)(x_1, \dots, x_n),
\end{aligned}$$

as needed. In particular, we have $dh_a = h_a d + (-1)^{n+1}(Id - *_a)$.

(ii) $dh'_a = d(h_a + *_a h_a + \dots + (*_a)^{k-1} h_a) = dh_a + (dh_a) * a + \dots + (dh_a) * a^{k-1} = h_a d + (-1)^{n+1}(Id - *_a) + (h_a d + (-1)^{n+1}(Id - *_a)) * a + \dots + (-1)^{n+1}(Id - *_a) * a^{k-1} = h_a d + (h_a * a) d + \dots + (h_a * a^{k-1}) d = h'_a d$.

(iii) It follows from the definition of rack, degenerate and quandle chain complex of X .

(iv) It suffices to consider the case, when there is $x \in X$, such that $b = a * x$ (the case $b = a \bar{*} x$ is equivalent to $a = b * x$ so there is no need to consider it separately).

Notice, that

$$\begin{aligned}
*_x(h'_a(w)) &= (h'_a(w)) * x = ((w + w * a + \dots + w * a^{k-1}), a) * x = \\
((w * x + (w * a) * x + \dots &+ (w * a^{k-1}) * x), b) = \\
((w * x + (w * x) * b + \dots &+ (w * x) * b^{k-1}), b) = \\
h'_b(w * x) &= h'_b(*_x(w)).
\end{aligned}$$

On the other hand, if w is a cycle then, by (i), $*_x(w)$ is a homologous cycle to w . Therefore, $h'_b(*_x(w))$ and $h'_b(w)$ are homologous by (ii). Similarly, $*_x(h'_a(w))$ and $h'_a(w)$ are homologous. Therefore, $h'_b(w)$ and $h'_a(w)$ are homologous.

Definition 9 (a) Let x_0 and x be two elements of a rack X . We say that x_0 is j -connected to x if there are elements $x_1, x_2, \dots, x_j$ such that $x_0 * x_1 * \dots * x_j = x$. Notice that if X is a quandle, then if x_0 is j -connected to x then it is also $(j+1)$ -connected to x (i.e., $x_0 * x_1 * \dots * x_j * x = x$). We also say that x is 0-connected to itself.

(b) We say that a rack X is j -connected with respect to x if for every $x_0 \in X$, x_0 is j (or less)-connected to x . We also define the distance $\rho(x_0, x)$ as the minimal j such that x_0 is j -connected to x . If such j does not exist, we write $\rho(x_0, x) = \infty$.

(c) We can visualize definitions (a) and (b) by constructing the oriented graph G_X (Cayley digraph of a rack with all elements as generators), whose vertices are elements of X , and edges (starting from x and ending at z) are triples (x, y, z) satisfying, $x * y = z$. We allow multiple edges and loops. In particular, if X is a quandle, any vertex x has a loop (x, x, x) .

(d) Recall that $\alpha : X \rightarrow X$ is an inner automorphism of length j if there are elements $x_1, x_2, \dots, x_j$ such that $\alpha(x) = x * x_1 * \dots * x_j$. We say that X is j -regular at $a \in X$ if for any pair of inner automorphisms of length j , α_1 and α_2 , such that $\alpha_1(x) = \alpha_2(x) = a$ for some x , we have $\alpha_1 = \alpha_2$. Notice that if X is a quandle, then a j -regular inner automorphism at a is also a $(j-1)$ -regular inner automorphism at a . Indeed, if $x_0 * x_1 * \dots * x_{j-1} = x_0 * y_1 * \dots * y_{j-1} = a$, then $x_0 * x_1 * \dots * x_{j-1} * a = x_0 * y_1 * \dots * y_{j-1} * a = a$, and by j -regularity, for any x we have $x * x_1 * \dots * x_{j-1} * a = x * y_1 * \dots * y_{j-1} * a = a$. Finally, $x * x_1 * \dots * x_{j-1} = x * y_1 * \dots * y_{j-1} = a$, showing $(j-1)$ -regularity.

Notice also, that when our j -regular condition is translated to Cayley digraphs, then it can be interpreted as “if two paths of length j starting at x_0 have the common endpoint, then the corresponding paths at any vertex have the same endpoint.”

(e) The rack X has the *quasigroup property*, if for any $a, b \in X$, the equation $a * x = b$ has exactly one solution. Notice that for a connected finite rack, the quasigroup property is equivalent to X being 1-regular.

Example 10 (1) Alexander quandles are j -regular for any j . To see this let $\alpha_1(x) = x * x_1 * \dots * x_j$ and $\alpha_2(x) = x * y_1 * \dots * y_j$. Then we have

$$\alpha_1(x) = (1-t)(x_j + tx_{j-1} + \dots + t^{j-1}x_1) + t^jx,$$

$$\alpha_2(x) = (1-t)(y_j + ty_{j-1} + \dots + t^{j-1}y_1) + t^jx.$$

Thus, $\alpha_1(x) - \alpha_2(x) = (1-t)((x_j - y_j) + t(x_{j-1} - y_{j-1}) + \dots + t^{j-1}(x_1 - y_1))$, which does not depend on x , proving j -regularity of Alexander quandles.

(2) Orbits of an Alexander quandle, A , are in bijection with the elements of

$A/(1-t)A$. Furthermore, every orbit of an Alexander quandle is 1-connected (and a quasigroup if the orbit is finite).

To see this, first notice that $x * y - x = (1-t)(y-x)$. Thus, elements in the same orbit are equal in $A/(1-t)A$. Conversely, if $y-x = (1-t)z$, then $y = x * (z+x)$. Thus y and x are in the same orbit and y is 1-connected to x ($\rho(x, y) = 1$), as required. Uniqueness of u with $x * u = y$ follows from the finiteness of the orbit (for an endomorphism of a finite set, an epimorphism is equivalent to a monomorphism). In the infinite case, an orbit is not necessary a quasigroup. This happens if $1-t$ annihilates some nonzero element of A .

In the following theorem, we generalize Theorem 9 of [21].

Theorem 11 *Let Q be a quandle consisting of s orbits with fixed points $\mathbf{a} = (a_1, a_2, \dots, a_s)$, with one a_i chosen from each orbit and satisfying the k -condition ($x * a_i^k = x$ for any x). We define a map $h'_{\mathbf{a}}: C_n^R(Q) \rightarrow C_{n+1}^R(Q)$ by $h'_{\mathbf{a}} = \sum_{i=1}^s h'_{a_i}$. It is a chain map as a sum of chain maps.*

- (1) *If Q is j -regular at each a_i , and each orbit of Q is a quasigroup, then there is a chain map $\bar{h}'_{\mathbf{a}}: C_{n+1}^R(Q) \rightarrow C_n^R(Q)$ such that on homology $(\bar{h}'_{\mathbf{a}} h'_{\mathbf{a}})_* = sk^2 Id$.*
- (2) *If each orbit of Q is j -connected to a_i , and Q is $(j+1)$ -regular at a_i , then there is a chain map $\bar{h}'_{\mathbf{a}}: C_{n+1}^R(Q) \rightarrow C_n^R(Q)$ such that $(\bar{h}'_{\mathbf{a}} h'_{\mathbf{a}})_* = sk^2 Id$ on homology.*
- (3) *In particular, if $H_n^R(Q)$ has no element of order dividing sk , then $(h'_{\mathbf{a}})_*: H_n^R(Q) \rightarrow H_{n+1}^R(Q)$ is a monomorphism.*

PROOF. We prove (2) of which (1) is a special case (that includes Alexander quandles).

Define $\bar{h}_{\mathbf{a}}(x_1, \dots, x_n, x_{n+1}) = (x_1, \dots, x_n) * u_1 * \dots * u_j$, where $x_{n+1} * u_1 * \dots * u_j = a_i$ for some i . The map is well defined because Q is j -regular at a_i .

In this situation, we define

$$\bar{h}'_{\mathbf{a}}(x_1, \dots, x_n, x_{n+1}) = \sum_{t=0}^{k-1} \bar{h}_{\mathbf{a}}(x_1, \dots, x_n, x_{n+1}) * a_i^t.$$

We prove that $\bar{h}'_{\mathbf{a}}: C_{n+1}^R(Q) \rightarrow C_n^R(Q)$ is a chain map. We have:

$$\begin{aligned} d\bar{h}'_{\mathbf{a}}(x_1, \dots, x_n, x_{n+1}) &= d\left(\sum_{t=0}^{k-1} (x_1, \dots, x_n) * u_1 * \dots * u_j * a_i^t\right) = \\ &= \sum_{t=0}^{k-1} d(x_1, \dots, x_n) * u_1 * \dots * u_j * a_i^t. \end{aligned}$$

On the other hand, we have:

$$\begin{aligned}
& \bar{h}'_{\mathbf{a}} d(x_1, \dots, x_n, x_{n+1}) = \\
& \bar{h}'_{\mathbf{a}}(d(x_1, \dots, x_n), x_{n+1}) + (-1)^{n+1} \bar{h}'_{\mathbf{a}}((x_1, \dots, x_n) - (x_1, \dots, x_n) * x_{n+1}) = \\
& \sum_{t=0}^{k-1} d(x_1, \dots, x_n) * u_1 * \dots * u_j * a_i^t \\
& + (-1)^{n+1} \bar{h}'_{\mathbf{a}}((x_1, \dots, x_n) - (x_1, \dots, x_n) * x_{n+1}).
\end{aligned}$$

Finally, we show that $(j+1)$ -regularity can be use to show that the last term is equal to zero. Namely, for some $a_{i'}$ depending on x_n , there are $y_1, \dots, y_j$ and $z_1, \dots, z_j$ such that we have $x_n * y_1 * \dots * y_j = a_{i'}$ (and so $x_n * y_1 * \dots * y_j * a_{i'} = a_{i'}$), and $x_n * x_{n+1} * z_1 * \dots * z_j = a_{i'}$. Thus by $(j+1)$ -regularity at $a_{i'}$, we have, for any x :

$$x * y_1 * \dots * y_j * a_{i'} = x * x_{n+1} * z_1 * \dots * z_j.$$

Thus,

$$\bar{h}'_{\mathbf{a}}((x_1, \dots, x_n) - (x_1, \dots, x_n) * x_{n+1}) = 0,$$

and $\bar{h}'_{\mathbf{a}}$ is a chain map.

Then, the formula $(\bar{h}'_{\mathbf{a}} h'_{\mathbf{a}})_* = sk^2 Id$ on homology is easy to check:

$$\begin{aligned}
& \bar{h}'_{\mathbf{a}} h'_{\mathbf{a}}(x_1, \dots, x_n) = \\
& \bar{h}'_{\mathbf{a}} \left(\sum_{j=0}^{k-1} \sum_{i=1}^s (x_1, \dots, x_n, a_i) * a_i^j \right) = \sum_{u=0}^{k-1} \left(\sum_{j=0}^{k-1} \sum_{i=1}^s (x_1, \dots, x_n) * a_i^j \right) * a_i^u = \\
& k \sum_{j=0}^{k-1} \sum_{i=1}^s (x_1, \dots, x_n) * a_i^j.
\end{aligned}$$

This holds for any chain, and if w is a cycle in $C_n^R(Q)$, then it is homologous to $w * a_i^j$. Therefore, $\bar{h}'_{\mathbf{a}} h'_{\mathbf{a}}(w) = k \sum_{j=0}^{k-1} \sum_{i=1}^s w * a_i^j$ is homologous to $sk^2 w$.

Corollary 12

- (i) If $(h'_{\mathbf{a}})_*$ is a non-trivial map (resp. monomorphism), and X is a homogeneous quandle (for any two elements x and y of X , there is an automorphism $X \rightarrow X$ sending x to y), then $(h'_a)_*$ is a nontrivial map (resp. monomorphism) for any $a \in X$.

- (ii) If A is a finite Alexander quandle, then $(\bar{h}'_a h'_a)_* = sk^2 Id$, where $s = |A/((1-t)A)|$ is the number of orbits of A , and k is the smallest number such that $(1-t^k)$ annihilates A .

PROOF. (i) If $(h'_a)_*(u) \neq 0$, then $(h'_{a_i})_*(u) \neq 0$ for some i , and thus, by homogeneity of X , $(h'_a)_*(x) \neq 0$ for some $x \in X$. The part (i) of Corollary 12 follows.

Part (ii) follows from (i), because any Alexander quandle is homogeneous. Recall that for any $a, b \in A$, the module automorphism $f_{a,b}: A \rightarrow A$ given by $f_{a,b}(x) = x + b - a$ is a quandle automorphism sending a to b . Also recall that because $x * a^k = x + (1-t^k)(b-x)$, A is a k -quandle if and only if $(1-t^k)$ annihilates A .

Example 13

- (i) $S_4 = \mathbb{Z}_2[t]/(1+t+t^2)$ is a connected Alexander 3-quandle with 4 elements, and $H_n^R(S_4)$ can have only 2-torsion. By Corollary 12, $(h'_a)_*: H_n^R(S_4) \rightarrow H_{n+1}^R(S_4)$ is a monomorphism. In particular, $\text{tor} H_n^R(S_4)$ is nontrivial (contains $\mathbb{Z}_2$) for $n \geq 2$; we have $H_2^R(S_4) = \mathbb{Z} \oplus \mathbb{Z}_2$.
- (ii) We computed that $H_2^Q(S_4) = \mathbb{Z}_2$, $H_3^Q(S_4) = \mathbb{Z}_2 \oplus \mathbb{Z}_4$, $H_4^Q(S_4) = \mathbb{Z}_2^2 \oplus \mathbb{Z}_4$, $H_5^Q(S_4) = \mathbb{Z}_2^5 \oplus \mathbb{Z}_4$, $H_6^Q(S_4) = \mathbb{Z}_2^9 \oplus \mathbb{Z}_4^2$, and $H_7^Q(S_4) = \mathbb{Z}_2^{17} \oplus \mathbb{Z}_4^3$. Generally, we conjecture that $\text{tor} H_n^Q(S_4) = \mathbb{Z}_2^{f'_n} \oplus \mathbb{Z}_4^{f_n}$, where $\{f_n\}$ are “delayed” Fibonacci numbers, that is, $f_n = f_{n-1} + f_{n-3}$, and $f(1) = f(2) = 0$, $f(3) = 1$; compare [21]. Furthermore, $g_n = f'_n + 2f_n = \log_2(|\text{tor} H_n^Q(S_4)|)$ satisfies $g_n = g_{n-1} + g_{n-2} + g_{n-4}$, with $g_1 = 0, g_2 = 1, g_3 = 3$, and $g_4 = 4$. According to this conjecture, we would have $H_8^Q(S_4) = \mathbb{Z}_2^{32} \oplus \mathbb{Z}_4^4$.
- (iii) Consider an Alexander quandle $A_{m,p(t)} = \mathbb{Z}_m[t^{\pm 1}]/(p(t))$, where $p(t)$ is a polynomial in variable t with the coefficients of the highest and lowest degree terms of t are invertible in $\mathbb{Z}_m$. $A_{m,p(t)}$ has $m^{\deg(p(t))}$ elements, and $s = \gcd(m, p(1))$ orbits. Then, for a connected quandle $A_{m,p(t)}$, the map $(h'_a)_*: H_n^R(A_{m,p(t)}) \rightarrow H_{n+1}^R(A_{m,p(t)})$ is a monomorphism. In particular, this holds for $p(x) = [k]_t = 1 + t + \dots + t^{k-1}$, with $\gcd(m, k) = 1$. Here, $A_{m,[k]_t}$ is a connected k -quandle (notice that $t^k - 1 = (t-1)[k]_t$ annihilates $A_{m,[k]_t}$).

3 Annihilation of rack homology

In this section, we offer an improvement of the results presented in [17,21], and we use it to show, in particular, that k^{n-2} annihilates $\text{tor} H_n(R_{2k})$ for k odd (Corollary 19). We conjecture, however, a much stronger result, at least for k prime:

Conjecture 14 *The number k annihilates $\text{tor}H_n(R_{2k})$, unless $k = 2^t$, $t > 1$. The number $2k$ is the smallest number annihilating $\text{tor}H_n(R_{2k})$ for $k = 2^t$, $t > 1$.*

We checked that $H_3^Q(R_8) = \mathbb{Z}^2 \oplus \mathbb{Z}_8^2$, $H_4^Q(R_8) = \mathbb{Z}^2 \oplus \mathbb{Z}_2^4 \oplus \mathbb{Z}_4^4 \oplus \mathbb{Z}_8^2$, and $H_3^Q(R_{16}) = \mathbb{Z}^2 \oplus \mathbb{Z}_{16}^2$.

To formulate our further results, we use definition of an X -set Y and the homology of the pair (X, Y) introduced in Definition 4.

We start from a fact, allowing as to detect torsion in H_4^Q of some quandles R_{4n} .

Consider two maps f, g defined as follows ([5, 21]):

$f: C_n^R(X, Y) \rightarrow C_{n-1}^R(X)$ given by $f(y, x_2, \dots, x_n) = (x_2, \dots, x_n)$, and
 $g: C_{n-1}^R(X) \rightarrow C_n^R(X, Y)$ given by $g(x_2, \dots, x_n) = \sum_{y \in Y} (y, x_2, \dots, x_n)$.

Proposition 15

(i) *The maps $(-1)^n f$, $(-1)^n g$, fg , and gf are chain maps. Furthermore, g sends degenerate elements in $C_{n-1}^R(X)$ to late degenerate elements in $C_n^R(X, Y)$.*

(ii) *$fg = |Y|Id$ on $C_{n-1}^R(X)$ and $gf(y, x_2, \dots, x_n) = \sum_{y \in Y} (y, x_2, \dots, x_n)$.*

PROOF. (i) We easily check that $df + fd = 0$ and $dg + gd = 0$. Thus, $(-1)^n f$, $(-1)^n g$, fg , and gf are chain maps.

(ii) It follows from a direct computation.

Corollary 16 *If $H_n^Q(R_{4k})$ has a $\mathbb{Z}_{4k}$ -torsion, then $H_{n+1}^Q(R_{4k})$ contains $\mathbb{Z}_2$ (compare Conjecture 14).*

PROOF. Let $X = R_{4k}$ and $Y = R_{4k}^{\text{even}}$ be an orbit composed of even numbers. Then $|Y| = 2k$ so if u is a generator of $\mathbb{Z}_{4k}$ in $H_n^R(R_{4k})$, then, by Corollary 16, $(gf)_*(u)$ is a nonzero element in $H_n^R(R_{4n})$. Therefore, $f_*(u)$ is a nontrivial element in $H_{n+1}^R(R_{4k})$. Because g_* is well defined on quandle homology, $g_*(u)$ is a nontrivial element in $H_{n+1}^Q(R_{4k})$ if u is a generator of $\mathbb{Z}_{4k}$ in $H_n^Q(R_{4k})$.

In the next theorem we analyze the chain map $gf: C_n^R(X, Y) \rightarrow C_n^R(X, Y)$, and show, in particular, that if Y is an orbit of X which is also a quasigroup, then gf is chain homotopic to $|Y|Id$ (compare [17, 4]).

Theorem 17 *Let (X, Y) be an X -set and let X_1 be a finite invariant subrack of X (e.g., an orbit of X). Consider the map $\phi_{X_1}: C_n^R(X, Y) \rightarrow C_n^R(X, Y)$ given by $\phi_{X_1}(y, x_2, \dots, x_n) = \sum_{x \in X_1} (y * x, x_2, \dots, x_n)$. Then*

- (1) ϕ_{X_1} is a chain map chain homotopic to $|X_1|Id$.
- (2) Let ${}_y*: X_1 \rightarrow Y$ be defined by ${}_y*(x) = y*x$, and assume that the cardinality of the set of elements of X_1 which send y to y' , for any pair $y, y' \in Y$, is finite and does not depend on y and y' . Let us denote this number by m so that $|X_1| = m|Y|$. Then $mgf = \phi_{X_1}$, which by (1) is chain homotopic to $|X_1|Id$.

In particular:

- (3) f_* and g_* are isomorphisms if homology is taken over any ring in which $|Q_1|$ is invertible, e.g., rational numbers.
- (4) If ${}_y*$ is always a bijection (e.g., $Y = X_1$, and X_1 is a quasigroup), then gf is chain homotopic to $|X_1|Id$.

PROOF. Consider a map $H = H_{X,Y,X_1}: C_n^R(X, Y) \rightarrow C_{n+1}^R(X, Y)$ given by

$$H(y, x_2, \dots, x_n) = (y, \sum_{x \in X_1} x, x_2, \dots, x_n).$$

It follows that

$$\begin{aligned} dH(y, x_2, \dots, x_n) = \\ |X_1|(y, x_2, \dots, x_n) - (y * (\sum_{x \in X_1} x), x_2, \dots, x_n) - Hd(y, x_2, \dots, x_n). \end{aligned}$$

Therefore, $dH + Hd = |X_1|Id - \phi_{X_1}$. We conclude that ϕ_{X_1} is a chain map chain homotopic to $|X_1|Id$. Parts (2), (3) and (4) follow straight from our conditions.

Corollary 18

- (i) Suppose that X_0 is an invariant subrack of X , with $|X_0|$ elements, that is a quasigroup. Then, if $\text{tor}H_n^R(X)$ is annihilated by N , then $\text{tor}H_{n+1}^R(X, X_0)$ is annihilated by $N|X_0|$.
- (ii) With the notation and assumptions as in Theorem 17(2), we have that if $\text{tor}H_n^R(X)$ is annihilated by N , then $\text{tor}H_{n+1}^R(X, Y)$ is annihilated by $N|X_1|/\gcd(N, m)$.

PROOF. (i) Let $a \in \text{tor}H_{n+1}^R(X, X_0)$. Then, by assumption, $Nf_*(a) = 0$ in $H_n^R(X)$. Therefore,

$$N|X_0|a = Ng_*f_*(a) = g_*(Nf_*(a)) = g_*(0) = 0.$$

(ii) Let $a \in \text{tor}H_{n+1}^R(X, Y)$. Then $Nf_*(a) = 0$ in $H_n^R(X)$. Thus,

$$\begin{aligned} 0 &= (m/(\gcd(N, m)))g_*(Nf_*(a)) = (N/(\gcd(N, m)(mg_*f_*(a))) = \\ &= (N/(\gcd(N, m))|X_1|Id, \end{aligned}$$

and $\text{tor}H_{n+1}^R(X, Y)$ is annihilated by $(N/(\gcd(N, m))|X_1|$.

Corollary 19 *For an even dihedral quandle R_{2k} , we have:*

- (i) $\text{tor}H_n^R(R_{2k})$ is annihilated by k^{n-2} , for k odd.
- (ii) $\text{tor}H_n^R(R_{2k})$ is annihilated by $2k^{n-1}$, for an even k .

PROOF. (i) If k is odd, then $Q_0 = Q_{\text{even}}$ composed of even numbers is an invariant quasigroup subquandle in Q . Thus, Theorem 17(4) applies. Furthermore,

$$H_n^R(R_{2k}) = H_n^R(R_{2k}, Q_0) \oplus H_n^R(R_{2k}, Q_{\text{odd}}),$$

where Q_{odd} is an invariant subquandle of R_{2k} composed of odd numbers. Also,

$$H_n^R(R_{2k}, Q_{\text{even}}) = H_n^R(R_{2k}, Q_{\text{odd}}),$$

as the quandle isomorphism $s_+ : R_{2k} \rightarrow R_{2k}$, given by $s_+(i) = i + 1$ (modulo $2k$), sends Q_{even} to Q_{odd} , and Q_{odd} to Q_{even} . Finally, by the result of Greene [13], $\text{tor}H_2^R(R_{2k}) = 0$ for k odd. The result follows by induction on n , starting from $n = 2$.

(ii) We use Theorem 17(2), for $Y = Q_{\text{even}}$ and $X_1 = R_{2k}$. Then, $m = |X_1|/|Y| = 2$, and $\gcd(m, N) = 2$, if N is even, and $\gcd(m, N) = 1$ if N is odd. The rest of the proof follows by induction on n , starting from $\text{tor}H_1(R_{2k}) = 0$.

Conjecture 20

- (i) $\text{tor}H_2^Q(R_{4k}) = \mathbb{Z}_2^2$.
- (ii) $H_n^Q(R_{2k})$ is annihilated by $2k^{n-2}$, for an even k .

We confirmed Conjecture 20(i) using GAP [12] for $k = 1, 2, 3, 4, 5, 6$. Part (ii) of Conjecture 20 follows from part (i) by Corollary 18(ii); compare Conjecture 14.

4 Partial derivatives and homological operations

By a homological operation of degree k we understand any homomorphism $H_*^R \rightarrow H_{*+k}^R$. A pre-homology operation of degree k is a chain map $h: C_*^R \rightarrow C_{*+k}^R$.

Below we give the necessary conditions for the map $h_w: C_*^R \rightarrow C_{*+\ell(w)}^R$ defined by $h_w(y, x_2, \dots, x_n) = (y, x_2, \dots, x_n, w)$, where $w \in C_{\ell(w)}^R(Q)$, to be a pre-homology operation of degree $\ell(w)$. To do this, we first define the partial derivatives $\frac{\partial^1}{\partial q}: C_n^R(Q) \rightarrow C_{n-1}^R(Q)$.

Definition 21 We will use the following standard notation.

$$\partial_i^0(x_1, \dots, x_n) = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n),$$

$$\partial_i^1(x_1, \dots, x_n) = (x_1 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n),$$

and

$$\partial^0 = \sum_{i=1}^n (-1)^i \partial_i^0, \quad \partial^1 = \sum_{i=1}^n (-1)^i \partial_i^1, \quad \partial = \partial^0 - \partial^1.$$

(i) For any $q \in Q$, we define:

$\frac{\partial^1}{\partial q}: C_n^R(Q) \rightarrow C_{n-1}^R(Q)$ by $\frac{\partial^1(x_1, \dots, x_n)}{\partial q} = \sum_{i=1}^n (-1)^i \partial_i^1 \delta_{x_i, q}$, where the Kronecker delta $\delta_{x_i, q} = 1$ if $x_i = q$, and 0 otherwise.

We have $\partial^1 = \sum_{q \in Q} \frac{\partial^1}{\partial q}$.

(ii) For any $q \in Q$, we define:

$\frac{\partial^0}{\partial q}: C_n^R(Q) \rightarrow C_{n-1}^R(Q)$ by $\frac{\partial^0(x_1, \dots, x_n)}{\partial q} = \sum_{i=1}^n (-1)^i \partial_i^0 \delta_{x_i, q}$.

We have $\partial^0 = \sum_{q \in Q} \frac{\partial^0}{\partial q}$.

(iii) $\frac{\partial^0}{\partial q}$ sends degenerate elements to degenerate elements for any rack Q . If Q is a quandle, then also $\frac{\partial^1}{\partial q}$ sends degenerate elements to degenerate elements.

Lemma 22 For a quandle Q , we have:

- (i) $\frac{\partial^0 \partial^0}{\partial q \partial q} = 0$,
- (ii) $\frac{\partial^0 \partial^0}{\partial q \partial q'} + \frac{\partial^0 \partial^0}{\partial q' \partial q} = 0$.
- (iii) $\frac{\partial^1 \partial^1}{\partial q \partial q} = 0$.
- (iv) $\partial^0 \frac{\partial^1}{\partial q} + \frac{\partial^1}{\partial q} \partial^0 = 0$.
- (v) $\frac{\partial^0 \partial^1}{\partial q \partial q} + \frac{\partial^1 \partial^0}{\partial q \partial q} = 0$

PROOF. Parts (i) and (ii) follow from a standard calculation, and no struc-

ture on Q is needed, it may be any set.

(iii) $\frac{\partial^1 \partial^1}{\partial q \partial q}(x_1, \dots, x_n) = \frac{\partial^1}{\partial q}(\sum_{i=1}^n (-1)^i (x_1 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n) \delta_{x_i, q}) = \sum_{j < i} (-1)^{i+j} ((x_1 * x_i)(x_j * x_i), \dots, (x_{j-1} * x_i)(x_j * x_i), x_{j+1} * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n) \delta_{x_i, q} \delta_{x_j * q, q} + \sum_{j > i} (-1)^{i+j-1} (x_1 * x_i * x_j, \dots, x_{i-1} * x_i * x_j, x_{i+1} * x_j, \dots, x_{j-1} * x_j, x_{j+1}, \dots, x_n) \delta_{x_i, q} \delta_{x_j, q} = 0$. We use in this calculation all axioms of quandle, in particular $q * q = q$ and $(x * q = q \Rightarrow x = q)$. In the language of face maps (see the next paragraph for a definition), we can write our property $\frac{\partial^1 \partial^1}{\partial q \partial q} = 0$ locally as:

$$\text{for } j < i: (\partial_j^1 \partial_i^1) \delta_{x_i, q} \delta_{x_j * q, q} = (\partial_{i-1}^1 \partial_j^1) \delta_{x_i, q} \delta_{x_j, q}.$$

Parts (iv) and (v):

Simplifying the notation from the proof of (iii), let us define the face map $\frac{\partial^1}{\partial q}(i)$ as $\partial_i^1 \delta_{x_i, q}$. That is,

$$\frac{\partial^1}{\partial q}(i)(x_1, \dots, x_n) = (x_1 * q, \dots, x_{i-1} * q, x_{i+1}, \dots, x_n)$$

if $x_i = q$, and 0 otherwise. Then we need to consider some cases. For $i < j$:

$$\begin{aligned} \partial_{j-1}^0 \frac{\partial^1}{\partial q}(i)(x_1, \dots, x_n) = \\ (-1)^{i+j-1} (x_1 * q, \dots, x_{i-1} * q, x_{i+1}, \dots, x_{j-1}, x_{j+1}, \dots, x_n) \end{aligned}$$

if $x_i = q$, and 0 otherwise.

For $i > j$:

$$\begin{aligned} \partial_j^0 \frac{\partial^1}{\partial q}(i)(x_1, \dots, x_n) = \\ (-1)^{i+j} (x_1 * q, \dots, x_{j-1} * q, x_{j+1} * q, x_{i-1} * q, x_{i+1}, \dots, x_n) \end{aligned}$$

if $x_i = q$, and 0 otherwise.

Similarly, for $i < j$ we have:

$$\begin{aligned} \frac{\partial^1}{\partial q}(i) \partial_j^0(x_1, \dots, x_n) = \\ (-1)^{i+j} (x_1 * q, \dots, x_{i-1} * q, x_{i+1}, \dots, x_{j-1}, x_{j+1}, \dots, x_n) \end{aligned}$$

if $x_i = q$, and 0 otherwise.

For $i > j$:

$$\begin{aligned} \frac{\partial^1}{\partial q}(i-1)\partial_j^0(x_1, \dots, x_n) = \\ (-1)^{i+j-1}(x_1 * q, \dots, x_{j-1} * q, x_{j+1} * q, x_{i-1} * q, x_{i+1}, \dots, x_n) \end{aligned}$$

if $x_i = q$, and 0 otherwise.

Therefore, for $i < j$:

$$\partial_{j-1}^0 \frac{\partial^1}{\partial q}(i) + \frac{\partial^1}{\partial q_i} \partial_j^0 = 0,$$

and for $i > j$:

$$\partial_j^0 \frac{\partial^1}{\partial q}(i) + \frac{\partial^1}{\partial q}(i-1)\partial_j^0 = 0.$$

Thus, for any rack it follows that

$$\partial^0 \frac{\partial^1}{\partial q} + \frac{\partial^1}{\partial q} \partial^0 = 0,$$

and for any quandle we have:

$$\frac{\partial^0}{\partial q} \frac{\partial^1}{\partial q} + \frac{\partial^1}{\partial q} \frac{\partial^0}{\partial q} = 0.$$

Remark 23 One could hope that $\frac{\partial^i \partial^j}{\partial q \partial q'} + \frac{\partial^j \partial^i}{\partial q' \partial q} = 0$ for $i, j \in \{0, 1\}$, $q, q' \in Q$. This is not the case as the following example illustrates:

Let $w = (q_0, q_1, q_2, q_0, q_1, q_2) \in C_6^Q(R_3)$. We have $\frac{\partial^1 w}{\partial q_1} = 0$, while

$$\frac{\partial^1 \partial^1 w}{\partial q_1 \partial q_2} = \frac{\partial^1(q_1, q_0, q_2, q_1, q_0)}{\partial q_1} = -(q_0, q_2, q_1, q_0) \neq 0.$$

4.1 Creating homology operations

Let $w \in C_\ell^R(Q)$. We define a map $h_w: C_n^R(Q) \rightarrow C_{n+\ell}^R(Q)$ by:

$$h_w(x_1, \dots, x_n) = (x_1, \dots, x_n, w).$$

Theorem 25 below gives the necessary conditions on w , so that $(h_w)_*$ is a homology operation, in the language reminiscent of searching for extrema

of a multi-variable function. Another criterion, applicable to non-connected quandles (e.g., R_{2k}), is given in Theorem 29.

Definition 24 We say that a chain $w \in C_\ell^R(Q)$ is an *extreme chain* if $\partial^0(w) = 0$, and all partial derivatives along w are equal to zero ($\frac{\partial^1 w}{\partial q} = 0$, for all q).

Theorem 25 Assume that $w \in C_\ell^R(Q)$ is an extreme chain. Then, h_w is a chain map, and $(h_w)_*$ is a homological operation. Furthermore, because h_w sends degenerate elements to degenerate elements, h_w is a chain map on quandle chains $C_n^Q(Q)$ ($\partial^0(w)$ and partial derivatives on quandle chains should be zero).

PROOF. We have

$$\begin{aligned} \partial h_w(u) &= \partial(u, w) = (\partial u, w) + (-1)^{n+1}((u, \partial^0(w)) - (\sum_{q \in Q} (u * q, \frac{\partial^1 w}{\partial q}))) = \\ h_w(\partial(u)) &+ (-1)^{n+1}((u, \partial^0(w)) - (\sum_{q \in Q} (u * q, \frac{\partial^1 w}{\partial q}))), \end{aligned}$$

and Theorem 25 follows.

Notice that our proof can be interpreted as demonstrating that the map $h_{\partial^0 w}$ is chain homotopic to the map $u \rightarrow (\sum_{q \in Q} (u * q, \frac{\partial^1 w}{\partial q}))$.

Example 26 Let $q_0, q_1, \dots, q_{N-1}$ be a sequence of “Fibonacci elements” of Q , that is, $q_i * q_{i+1} = q_{i+2}$, with indices taken modulo N . Then:

- (i) [21] $s(q_0, q_1) = \sum_{i=0}^{N-1} (q_i, q_{i+1})$ satisfies the conditions of Theorem 25.
- (ii) $w = \partial^0 \frac{\partial^1(q_0, q_1, \dots, q_{N-1}, q_0)}{\partial q_0}$ satisfies the conditions of Theorem 25 for a quandle chain complex.

Example 27

- (i) The chain

$$w = -(2, 4, 1) - (3, 2, 1) - (4, 3, 1) + (1, 2, 4) + (1, 3, 2) + (1, 4, 3),$$

where $1, 2, 3, 4$ denote the elements $0, 1, t$, and $1 + t$ of the quandle $S_4 = \mathbb{Z}_2[t]/(t^2 + t + 1)$, is an extreme chain, and for $g = (1, 2) + (2, 4) + (4, 1)$, (g, w) gives $\mathbb{Z}_2$ in $H_5^Q(S_4)$.

- (ii) For a dihedral quandle R_3 , the chain

$$w = -(1, 0, 1, 2, 0) - (1, 2, 0, 2, 0) - (2, 0, 2, 1, 0) - (2, 1, 0, 1, 0)$$

$$+(0, 1, 0, 1, 2) + (0, 1, 2, 0, 2) + (0, 2, 0, 2, 1) + (0, 2, 1, 0, 1)$$

represents $\mathbb{Z}_3$ in $H_5^Q(R_3)$. It also satisfies the conditions of the Theorem 25, and the operation h_w applied to low dimensional cycles gives torsion elements in homology groups of higher degrees.

Remark 28 In [20] we introduced the concept of the n -th Burnside kei, that is a kei (involutive quandle) Q in which any pair of elements, a_0, a_1 , satisfies the relation $a_n(a_0, a_1) = a_0$. Here we use the notation

$$a_n = a_n(a_0, a_1) = \underbrace{\dots a_1 * a_0 * a_1 * a_0 * a_1}_{\text{total of n letters}}.$$

Then the Fibonacci condition $a_{i+2} = a_i * a_{i+1}$ is satisfied, and we can define the Fibonacci element $s(a_0, a_1) = \sum_{i=0}^{n-1} (a_i, a_{i+1})$ which is an extreme chain in $C_2^R(Q)$, so it can be used to define a homology operation $h_{s(a_0, a_1)}$. In fact, the assumption that Q is a kei is not needed here, so we define the n -th Burnside quandle to be the quandle for which the equation $a_n(a_0, a_1) = a_0$ holds for any pair of elements, a_0, a_1 . In this setting, $s(a_0, a_1)$ is also an extreme chain and $h_{s(a_0, a_1)}$ is a homology operation; compare Example 38.

Theorem 29 *Let Q be a quandle and Q_1 its invariant subquandle, such that $*_q : Q \rightarrow Q$ does not depend on the choice of $q \in Q_1$. Assume also that (any) $q \in Q_1$ is a k -element of Q (i.e., $x * q^k = x$, for any $x \in Q$). Then $h'_w = \sum_{i=0}^{k-1} h_w * q^i$ is a chain map for any $w \in \mathbb{Z}Q_1^\ell$.*

PROOF. We have

$$\begin{aligned} \partial h'_w(x_1, \dots, x_n) &= \partial \sum_{i=0}^{k-1} h_w * q^i = h'_w \partial(x_1, \dots, x_n) \\ &+ (-1)^{n+1} \sum_{i=0}^{k-1} ((x_1, \dots, x_n, \partial^0 w) - ((x_1, \dots, x_n) * q, \partial^1 w)) * q^i. \end{aligned}$$

Theorem 29 follows, as Q_1 is a trivial quandle with $\partial^0 w = \partial^1 w$.

Example 30 The dihedral quandle R_4 is an example of a quandle which motivated Theorem 29 (here, $Q_1 = \{0, 2\}$). We generalize it as follows. Consider two trivial quandles: Q_0 equal to $\mathbb{Z}_{k_0}$ as a set, and Q_1 equal to $\mathbb{Z}_{k_1}$ as a set. We define the quandle Q_{k_0, k_1} on the set $Q_0 \cup Q_1$ by $a * b = a + 1$ for $a \in Q_0$ and $b \in Q_1$ or $a \in Q_1$ and $b \in Q_0$. Our quandle Q (with Q_1 or Q_0 as invariant subquandle) satisfies the conditions of Theorem 29. For example, we have three such quandles of size 6 (Q_{5+1} , Q_{4+2} , Q_{3+3}). Notice that $Q_{2,2}$ is the quandle R_4 , and the quandle Q_{k_0, k_1} is an $lcm(k_0, k_1)$ -quandle.

4.2 Naturality of homological operations

The naturality we have in mind is defined as follows. Suppose that we have a pre-homological operation for a given rack X_1 , that is, a chain map $h: C_n^R(X_1) \rightarrow C_{n+k}^R(X_1)$, and there is a rack homomorphism $f: X_1 \rightarrow X_2$. Then, there is a uniquely defined pre-homological operation $T(h): C_n^R(X_2) \rightarrow C_{n+k}^R(X_2)$, such that $f_{\#}h = T(h)f_{\#}$, where $f_{\#}: C_*^R(X_1) \rightarrow C_*^R(X_2)$ is the chain map induced by f (we could also go straight to homology and write $f_*h = (T(h))_*f_*$ as a condition for a homological operation).

Let $f: Q_1 \rightarrow Q_2$ be a quandle homomorphism, and $f_*: H_n^W(Q_1) \rightarrow H_n^W(Q_2)$ be the induced homomorphism, where $W = R, Q$ or D . We define $f^{(n)}: Q_1^n \rightarrow Q_2^n$ by

$$f^{(n)}(x_1, \dots, x_n) = (f(x_1), \dots, f(x_n)),$$

and extend this map linearly to the group homomorphism $f^{(n)}: \mathbb{Z}Q_1^n \rightarrow \mathbb{Z}Q_2^n$.

Theorem 31

- (i) If a is a k -element in Q_1 , and $f(a)$ is a k -element in Q_2 , then $fh'_a = h'_{f(a)}f$, so h'_a is natural.
- (ii) If $w \in C_\ell^R(Q_1)$ is an extreme chain (as in Theorem 25), then

$$(h_{f^{(\ell)}(w)})_*: H_n^R(Q_2) \rightarrow H_{n+\ell}^R(Q_2)$$

is a homological operation.

- (iii) The operation h_w is natural.

PROOF. (i) We have

$$fh'_a(x_1, \dots, x_n) = f\left(\sum_{i=0}^{k-1} (*_a)^i(x_1, \dots, x_n, a)\right) =$$

$$\sum_{i=0}^{k-1} (*_{f(a)})^i(f(x_1), \dots, f(x_k), f(a)) = h'_{f(a)}f,$$

as needed.

- (ii) Consider $\frac{\partial^1 f^{(\ell)}(w)}{\partial p}$. If p is not in the image $f(Q_1)$, then p is not an element occurring in $f^{(\ell)}(w)$, so $\frac{\partial^1 f^{(\ell)}(w)}{\partial p} = 0$. Let $q_1, q_2, \dots, q_k$ be the elements in the

preimage $f^{-1}(p)$ that are in w . We will show that

$$\frac{\partial^1 f^{(\ell)}(w)}{\partial p} = f^{(\ell-1)} \sum_{i=1}^k \frac{\partial^1 w}{\partial q_i}.$$

It is enough to show this formula in the case $w = (x_1, \dots, x_\ell)$, where $x_i \in Q_1$ for $i = 1, \dots, \ell$:

$$\begin{aligned} f^{(\ell-1)} \sum_{i=1}^k \frac{\partial^1(x_1, \dots, x_\ell)}{\partial q_i} &= \\ f^{(\ell-1)} \sum_{i=1}^k \sum_{j=1}^{\ell} (-1)^j (x_1 * q_i, \dots, x_{j-1} * q_i, x_{j+1}, \dots, x_\ell) \delta_{x_j, q_i} &= \\ \sum_{i=1}^k \sum_{j=1}^{\ell} (-1)^j f^{(\ell-1)}(x_1 * q_i, \dots, x_{j-1} * q_i, x_{j+1}, \dots, x_\ell) \delta_{x_j, q_i} &= \\ \sum_{i=1}^k \sum_{j=1}^{\ell} (-1)^j (f(x_1) * f(q_i), \dots, f(x_{j-1}) * f(q_i), f(x_{j+1}), \dots, f(x_\ell)) \delta_{x_j, q_i} &= \\ \sum_{j=1}^{\ell} (-1)^j (f(x_1) * p, \dots, f(x_{j-1}) * p, f(x_{j+1}), \dots, f(x_\ell)) \delta_{f(x_j), p} &= \frac{\partial^1 f^{(\ell)}(w)}{\partial p}. \end{aligned}$$

Similarly, if $\partial^0 = 0$, then $\partial^0 f^{(\ell)}(w)$.

(iii) We have to check that $f^{(n+\ell)} h_w = h_{f^{(\ell)}(w)} f^{(n)}$. Indeed,

$$\begin{aligned} f^{(n+\ell)} h_w(x_1, \dots, x_n) &= f^{(n+\ell)}(x_1, \dots, x_n, w) = (f(x_1), \dots, f(x_n), f(w)) = \\ h_{f^{(\ell)}(w)}(f(x_1), \dots, f(x_n)) &= h_{f^{(\ell)}(w)} f^{(n)}(x_1, \dots, x_n), \end{aligned}$$

as needed.

Remark 32 Carter, Elhamdadi, and Saito [2] proposed a generalization of quandle homology to twisted quandle homology with $\partial^T = t\partial^0 - \partial^1$. Any extreme chain $w \in C_\ell^R(Q)$ is also an extreme chain in the twisted homology (or any homology with the differential of the form $a\partial^0 \pm b\partial^1$), and can be used to construct a natural homology operation. This awaits detailed exploration.

5 Further applications

The techniques we have built so far can be used for some concrete calculations of quandle homology. We will illustrate it by several examples for both finite and infinite quandles. We start from a simple lemma crucial for our examples.

Lemma 33 *Consider a rack X and its subrack X_0 , with an embedding $i: X_0 \rightarrow X$. We say that a rack epimorphism $r: X \rightarrow X_0$ is a rack twist-retraction¹ if $ri \in \text{Aut}(X_0)$, and a rack retraction if $ri = \text{Id}_{X_0}$. Then $i_*: H_n^R(X_0) \rightarrow H_n^R(X)$ is a monomorphism, and $r_*: H_n^R(X) \rightarrow H_n^R(X_0)$ is an epimorphism.*

If X is an Alexander quandle, then the retraction corresponds to an $\mathbb{Z}[t^{\pm 1}]$ -module epimorphism which splits:

Corollary 34 *Consider a $\mathbb{Z}[t^{\pm 1}]$ -module $A = A_1 \oplus A_2$ with embeddings $i_1: A_1 \rightarrow A_1 \oplus A_2$, $i_2: A_2 \rightarrow A_1 \oplus A_2$, and projections $r_1: A_1 \oplus A_2 \rightarrow A_1$, $r_2: A_1 \oplus A_2 \rightarrow A_2$. Naturally, $r_1 i_1 = \text{Id} = r_2 i_2$, so r_1 and r_2 are retractions. We have:*

- (i) $(i_1)_*: H_n^W(A_1) \rightarrow H_n^W(A)$, $(i_2)_*: H_n^W(A_2) \rightarrow H_n^W(A)$
are monomorphisms.
- (ii) If $\gcd(|A_1|, |A_2|) = 1$, then

$$H_n^W(A_1) \oplus H_n^W(A_2)$$

embeds in $H_n^W(A)$.

PROOF. The first fact follows from Lemma 33. The second conclusion follows from the fact that $|A_i|^n$ annihilates $H_n^W(A_i)$, for $i = 1, 2$.

Corollary 35

- (i) If $\gcd(m, k) = 1$, then $\mathbb{Z}_{mk} = \mathbb{Z}_m \oplus \mathbb{Z}_k$ and

$$\text{tor} H_n^W(R_m) \oplus \text{tor} H_n^W(R_k) \subset H_n^W(R_{mk}).$$

- (ii) For k odd,

$$\text{tor} H_n^W(R_k) \oplus \text{tor} H_n^W(R_k) \subset H_n^W(R_{2k}).$$

¹ Of course it is always possible, having a twist-retraction r , to consider a retraction $r' = (ri)^{-1}r$ (we have $r'i = (ri)^{-1}ri = \text{Id}_{X_0}$). However, it is sometimes convenient to work with twist-retractions of racks.

(iii) For k odd, prime, such that $\gcd(m, k) = 1$, $H_3^Q(R_{mk})$ and $H_4^Q(R_{mk})$, contain $\mathbb{Z}_k$ (compare Theorem 7 (ii)).

Inequality in Corollary 35(ii) is seldom an equality, even for k odd prime. For example, $H_4^Q(R_6) = \mathbb{Z}_3^6 = H_4^Q(R_3) \oplus H_4^Q(R_3) \oplus \mathbb{Z}_3^4$. The reason may be that R_6 , not being connected, allows for more homological operations than R_3 (e.g., two different h'_a operations). We conjecture, however, the following:

Conjecture 36

$$(i) \quad \text{tor}(H_{2n}^Q(R_4)) = (\text{tor}(H_{2n-1}^Q(R_4)))^2 \oplus \mathbb{Z}_2^2 = \mathbb{Z}_2^{2(4^{n-1}-1)/3}$$

$$\text{tor}(H_{2n+1}^Q(R_4)) = (\text{tor}(H_{2n}^Q(R_4)))^2 = \mathbb{Z}_2^{4(4^{n-1}-1)/3}.$$

(ii) For k odd:

$$\text{tor}(H_{2n}^Q(R_{2k})) = (\text{tor}(H_{2n-1}^Q(R_{2k})))^2 \oplus \mathbb{Z}_k^2 = \mathbb{Z}_k^{(5 \cdot 4^{n-1} - 2)/3}, \text{ for } n > 1,$$

$$\text{tor}(H_{2n+1}^Q(R_{2k})) = (\text{tor}(H_{2n}^Q(R_{2k})))^2 = \mathbb{Z}_k^{2(5 \cdot 4^{n-1} - 2)/3}.$$

$$(iii) \quad 2^{k-1} \text{tor}(H_n^Q(R_{2k})) = \mathbb{Z}_2^{2f_n}, \text{ for } k > 1.$$

Here, $\{f_n\}$ denotes the sequence of “delayed” Fibonacci numbers, as in Example 2.6.

Example 37 Corollary 34 can be applied directly to an interesting family of Alexander quandles $A_{p, [k_1]_t [k_2]_t} = \mathbb{Z}_p[t^{\pm 1}] / ([k_1]_t \cdot [k_2]_t)$, with $\gcd(k_1, k_2) = 1$. Recall the notation $[k]_t = 1 + t + \dots + t^{k-1}$, $A_{p, [k]_t} = \mathbb{Z}_p[t^{\pm 1}] / ([k]_t)$. $A_{p, [k_1]_t [k_2]_t}$ splits to $A_{p, [k_1]_t} \oplus A_{p, [k_2]_t}$, so

$$\text{tor} H_n^W(A_{p, [k_1]_t}) \subset \text{tor} H_n^W(A_{p, [k_1]_t [k_2]_t}).$$

For example, $A_{p, [2]_t} = R_p$, and $A_{2, [3]_t} = S_4$, so we know a lot about their quandle homology. We also computed, for instance, that $H_2^Q(A_{2, [5]_t}) = \mathbb{Z}_2^2$, and $H_2^Q(A_{2, [4]_t}) = H_2^Q(A_{2, [6]_t}) = \mathbb{Z}^2 \oplus \mathbb{Z}_4$.

Example 38 Consider the Fibonacci quandle $F_n = \{a_0, a_1, \dots, a_{n-1} \mid a_2 = a_0 * a_1, a_3 = a_2 * a_1, \dots, a_{n-1} = a_{n-2} * a_{n-3}, a_0 = a_{n-2} * a_{n-1}, a_1 = a_{n-1} * a_0\}$. Then we have a Fibonacci element $s(a_0, a_1) = \sum_{i=0}^{n-1} (a_i, a_{i+1})$ which can be used for a homological operation $h_{s(a_0, a_1)}$. We also have an epimorphism $r: F_n \rightarrow R_n$ given by $r(a_j) = j$. This can be used to produce nontrivial elements in $H_k^W(F_n)$. Notice that F_n is the fundamental quandle of the torus link of type $(2, n)$, $T_{2, n}$; compare Figure 1.

(i) $H_n^Q(F_4)$ contains the free part for $2 \leq n \leq 7$, because we found that in $H_n^Q(R_4)$ the elements:




Fig. 1. The torus link $T_{2,n}$ with Fibonacci quandle labeling

$s(0, 1)$, $h_{s(0,1)}((0))$, $h_{s(0,1)}(s(0, 1))$, $h_{s(0,1)}h_{s(0,1)}((0))$, $h_{s(0,1)}h_{s(0,1)}(s(0, 1))$, and $h_{s(0,1)}h_{s(0,1)}h_{s(0,1)}((0))$ represent $\mathbb{Z}$. Compare [9], where it was shown that $H_2^Q(F_4) = \mathbb{Z}^2$.

- (ii) $H_3^Q(F_p)$ is nontrivial for p odd prime. It follows from the fact that $r_*((a_0, s(a_0, a_1)))$ is a non-zero element in $H_3^Q(R_p) = \mathbb{Z}_p$. In fact, in all cases we checked $h_{s(0,1)}: C_n^Q(R_k) \rightarrow C_{n+2}^Q(R_k)$ induces a monomorphism on homology [21]. From this follows that $H_{2n+1}^Q(F_3)$ is nontrivial for $n \leq 5$. One can conjecture that $h_{s(a_0, a_1)}: C_n^Q(F_k) \rightarrow C_{n+2}^Q(F_k)$ induces a monomorphism on homology. Our methods are not yet sufficient to decide whether $H_n^Q(F_k)$ can have torsion elements. To answer this question, we plan to use the detailed description of F_3 in [22] (the fundamental quandle of the trefoil knot was interpreted there as a symplectic quandle).

We end with a few rather general conjectures (39, 41, 42), and two new examples illustrating them:

Conjecture 39 *Let $r: Q_1 \rightarrow Q_2$ be an epimorphism of finite connected quandles, such that $|Q_1| = k|Q_2|$, and $\gcd(k, |Q_2|) = 1$. Then, $r_*: H_n^W(Q_1) \rightarrow H_n^W(Q_2)$ is an epimorphism, and if restricted to the part of $H_n^R(Q_1)$ not annihilated by k^j for some j , it is an isomorphism.*

Example 40 Consider the 6-element connected quandles Q_1 and Q_2 which are the last two quandles from the page 195 of [6]. We have an epimorphism $r_1: Q_1 \rightarrow R_3$, and $r_2: Q_2 \rightarrow R_3$. As far as computation of homology was performed, Conjecture 39 holds. In particular, non-2 torsion part of homology of Q_i , $i = 1, 2$, is the same as that for R_3 .

For example, Q_2 , the last quandle in [6], is a 4-quandle, given by the following

*-multiplication table:

*	1	2	3	4	5	6
1	1	1	6	3	4	5
2	2	2	4	5	6	3
3	4	6	3	3	3	1
4	5	3	1	4	2	4
5	6	4	5	1	5	2
6	3	5	2	6	1	6

and $r_2: Q_2 \rightarrow R_3$ is given by $r_2(1) = r_2(2) = 0$, $r_2(3) = r_2(5) = 1$, and $r_2(4) = r_2(6) = 2$. $s(1, 3) = (1, 3) + (3, 6) + (6, 1)$ is a Fibonacci chain in $C_2^Q(Q_2)$, with $r_2(s(1, 3)) = s(0, 1) \in C_2^Q(R_3)$.

If we combine Example 40 with the prediction that $3 \cdot 2^i$ annihilates torsion in homology of Q_1 and Q_2 , then we can suggest:

Conjecture 41 *Let $oddtor$ denote the odd part of the torsion of a group. Then*

$$oddtor(H_n^W(Q_1)) = oddtor(H_n^W(Q_2)) = oddtor(H_n^W(R_3)) = \mathbb{Z}_3^{f_n},$$

where f_n is as defined before.

The last, rather general conjecture should be the first step to understand torsion in homology of quandles, but, until now, the only nontrivial quandle for which it is solved is the dihedral quandle R_3 [21].

Conjecture 42 *If Q is a finite quasigroup quandle, then $|Q|$ annihilates torsion of its homology.*

Notice that *quasigroup* cannot be replaced by *connected* as 6 does not annihilate the torsion of homology of Q_2 . In particular, $H_3^Q(Q_2) = \mathbb{Z}_8 \oplus \mathbb{Z}_3$ (see [7]).

6 Acknowledgements

M. Niebrzydowski was partially supported by the Louisiana Board of Regents grant (# LEQSF(2008-11)-RD-A-30), and by the research award from the University of Louisiana at Lafayette.

J. H. Przytycki was partially supported by the NSA grant (# H98230-08-1-0033), by the NSF grant (# 0745204), and by the CCAS/UFF award.

We would like to thank the referee and T. Nosaka for many suggestions that improved our paper.

References

- [1] N. Andruskiewitsch, M. Graña, From racks to pointed Hopf algebras, *Adv. Math.* 178 (2003) 177-243.
- [2] S. Carter, M. Elhamdadi, M. Saito, Twisted quandle homology theory and cocycle knot invariants, *Algebraic and Geometric Topology* 2 (2002) 95-135.
- [3] S. Carter, D. Jelsovsky, S. Kamada, L. Langford, M. Saito, State-sum invariants of knotted curves and surfaces from quandle cohomology, *Electron. Res. Announc. Amer. Math. Soc.* 5 (1999) 146-156.
- [4] S. Carter, D. Jelsovsky, S. Kamada, M. Saito, Quandle homology groups, their Betti numbers, and virtual knots, *J. Pure Appl. Algebra* 157 (2001) 135-155.
- [5] S. Carter, D. Jelsovsky, S. Kamada, M. Saito, Shifting homomorphisms in quandle cohomology and skeins of cocycle knot invariants, *Journal of Knot Theory and its Ramifications* 10 (2001) 579-596.
- [6] S. Carter, S. Kamada, M. Saito, Surfaces in 4-space, *Encyclopaedia of Mathematical Sciences, Low-Dimensional Topology III*, R.V.Gamkrelidze, V.A.Vassiliev, Eds., Springer-Verlag Berlin-Heidelberg, 2006, 213pp.
- [7] S. Carter, S. Kamada, M. Saito, Geometric interpretations of quandle homology, *Journal of Knot Theory and its Ramifications* 10 (2001) 345-386.
- [8] P. Etingof, M. Graña, On rack cohomology, *J. Pure Appl. Algebra* 177 (2003) 49-59.
- [9] M. Eisermann, Homological characterization of the unknot, *J. Pure Appl. Algebra* 177 (2003) 131-157.
- [10] R. Fenn, C. Rourke, Racks and links in codimension two, *Journal of Knot Theory and its Ramifications* 1 (1992) 343-406.
- [11] R. Fenn, C. Rourke and B. Sanderson, James bundles and applications, preprint. e-print: <http://www.maths.warwick.ac.uk/~cpr/ftp/james.ps>
- [12] The GAP Group, GAP – Groups, Algorithms, and Programming, <http://www.gap-system.org>.
- [13] M. Greene, Some results in geometric topology and geometry, Ph.D. thesis, University of Warwick, advisor: Brian Sanderson, 1997.

- [14] D. Joyce, An algebraic approach to symmetry with applications to knot theory, Ph.D. thesis, University of Pennsylvania, advisor: Peter Freyd, 1979.
- [15] D. Joyce, A classifying invariant of knots: the knot quandle, *J. Pure Appl. Algebra* 23 (1982) 37-65.
- [16] S. Kamada, Quandles with good involutions, their homologies and knot invariants, in: *Intelligence of Low Dimensional Topology 2006*, Eds. J. S. Carter et. al., World Scientific Publishing Co., 2007, pp. 101-108.
- [17] R. A. Litherland, S. Nelson, The Betti numbers of some finite racks, *J. Pure Appl. Algebra* 178 (2003) 187-202.
- [18] S. Matveev, Distributive groupoids in knot theory, *Math. Sbornik* 119 (1982) 78-88. Eng. transl.: *Math. USSR Sbornik* 47 (1984) 73-83.
- [19] T. Mochizuki, Some calculations of cohomology groups of finite Alexander quandles, *J. Pure Appl. Algebra* 179 (2003) 287-330.
- [20] M. Niebrzydowski, J. H. Przytycki, Burnside Kei, *Fundamenta Mathematicae* 190 (2006) 211-229.
- [21] M. Niebrzydowski, J. H. Przytycki, Homology of dihedral quandles, *J. Pure Appl. Algebra* 213 (2009) 742-755.
- [22] M. Niebrzydowski, J. H. Przytycki, The quandle of the trefoil as the Dehn quandle of the torus, *Osaka Journal of Mathematics* 46 (2009) 645-659.
- [23] T. Ohtsuki, Quandles, in: *Problems on invariants of knots and 3-manifolds, Geometry and Topology Monographs* 4 (2003) 455-465.
- [24] M. Takasaki, Abstraction of symmetric transformation, (in Japanese) *Tohoku Math. J.* 49 (1942/3) 145-207.